Gastroenterology

SUPREP Colonoscopy Prep

Procedure Before Noon

Page 1 of 1

- Purchase from the Pharmacy
 - One (1) box of Simethicone (Gas-X) Extra-Strength 125mg oral soft gel capsules (over the counter)
 - SUPREP Solution (prescribed pick up from pharmacist)

Bowel Preparation Instructions:

The day before your colonoscopy, diet is to consist of CLEAR LIQUIDS ONLY (no solid food).

• Do not consume liquids that are red or purple in color. Do not consume milk products or juices with pulp. Clear liquids include: water, black coffee, tea, apple juice, white grape or cranberry juice, lemonade, sports drinks (Gatorade, PowerAde, etc.), clear beef, chicken or vegetable broth, Jell-O, popsicles, frozen sorbet, or Italian ice. Alcoholic beverages are not permitted.

SUPREP Bowel Prep solution can be refrigerated.

- 1. **At 5:00 PM,** pour one (1) 6-ounce bottle of SUPREP liquid into the mixing container. Add cold drinking water or any clear liquid to the 16-ounce line on the container and mix. Drink all the liquid in the container. *You MUST drink TWO (2) more 16-ounce containers of water or other clear liquid following the PREP solution.* You can continue to drink clear liquids during the evening.
- 2. At 10:00 PM pour the last 6-ounce bottle of SUPREP liquid into the mixing container. Add cold drinking water or any other clear liquid to the 16-ounce line on the container and mix. Drink all the liquid in the container. You MUST drink TWO (2) more 16-ounce containers of water or other clear liquid following the tablets. At the same time, take two (2) extra strength Simethicone (Gas-X) oral soft gel capsules (250mg total).

You MUST finish this process by 12:00AM.

⇒ After midnight, consume nothing further by mouth except a sip of water for necessary medications.

Contact the on-call Gastroenterologist **(941) 262-0400** if any problems arise during the bowel preparation process.

Prep Guide

Dark and murky
Brown and murky
Dark orange and semi-clear
Light orange and mostly clear
Yellow and clear

