Gastroenterology

SUFLAVE Colonoscopy Prep Procedure After Noon

Page 1 of 1

Purchase from the Pharmacy

- One (1) box of Simethicone (Gas-X) Extra-Strength 125mg oral soft gel capsules (over the counter)
- SUFLAVE Solution (prescribed pick up from pharmacist)

Bowel Preparation Instructions

The day before your colonoscopy, diet is to consist of CLEAR LIQUIDS ONLY (no solid food).

- Do not consume liquids that are red or purple in color. Do not consume milk products or juices
 with pulp. Clear liquids include water, black coffee, tea, apple juice, white grape or cranberry
 juice, lemonade, sports drinks (Gatorade, PowerAde, etc.), clear beef, chicken or vegetable
 broth, Jell-O, popsicles, frozen sorbet, or Italian ice. Alcoholic beverages are not permitted.
- SUFLAVE bowel prep solution can be refrigerated and must be consumed within 24 hours of being mixed.

• The day before your procedure, at 8:00AM:

 Prepare the SUFLAVE prep solution according to the package instructions included in your prep kit.

The day before your procedure, at 7:00PM:

 Consume one (1) 8-ounce glass of prep mixture every 20 minutes until the entire first bottle of prep solution has been consumed.
 This should be followed by one (1) 16-ounce cup of clear liquid.

The morning of your procedure, at 5:00AM:

- Consume one (1) 8-ounce glass of prep mixture every 20 minutes until the entire second bottle of prep solution has been consumed. This should be followed by one (1) 16-ounce cup of clear liquid.
- Take two (2) extra strength Simethicone (Gas-X) oral soft gel capsules (250mg total).
- ⇒ You may continue to consume clear liquids until four (4) hours before your procedure time. In all circumstances, consume nothing further by mouth within four (4) hours of your procedure time except for a sip of water for necessary medications.

Contact the on-call Gastroenterologist at **(941) 262-0400** if any problems arise during the bowel preparation process.



Prep Guide

